

University of Groningen

Development of macronutrient composition of very preterm human milk

Maas, Yolande G. H.; Gerritsen, Jeanet; Hart, Augustinus A. M.; Hadders-Algra, Mijna; Ruijters, Jan M.; Tamminga, Pieter; Mirmiran, Majid; Spekrijse, Henk

Published in:
British Journal of Nutrition

DOI:
[10.1017/S0007114598001743](https://doi.org/10.1017/S0007114598001743)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Maas, Y. G. H., Gerritsen, J., Hart, A. A. M., Hadders-Algra, M., Ruijters, J. M., Tamminga, P., Mirmiran, M., & Spekrijse, H. (1998). Development of macronutrient composition of very preterm human milk. *British Journal of Nutrition*, 80(1), 35-40. <https://doi.org/10.1017/S0007114598001743>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Development of macronutrient composition of very preterm human milk

Yolanda G. H. Maas^{1*}, Jeanet Gerritsen¹, Augustinus A. M. Hart², Mijna Hadders-Algra³,
Jan M. Ruijter⁴, Pieter Tamminga¹, Majid Mirmiran⁵ and Henk Spekrijse⁴

¹Department of Neonatology and ²Department of Clinical Epidemiology and Biostatistics, Academic Medical Center, University of Amsterdam, Emma Childrens' Hospital, PO Box 22700, 1100 DE Amsterdam, The Netherlands

³Department of Medical Physiology, University of Groningen, Bloemsingel 10, 9712 KZ Groningen, The Netherlands

⁴The Netherlands Ophthalmic Research Institute and Laboratory of Medical Physics, PO Box 12141, 1100 AC Amsterdam, The Netherlands

⁵Netherlands Institute for Brain Research, Meibergdreef 33, 1105 AZ Amsterdam, The Netherlands

(Received 24 July 1997 – Revised 8 January 1998 – Accepted 27 January 1998)

The effects of gestational age at delivery (GA), postnatal age (PNA) and post-menstrual age (PMA = PNA + GA, an indicator of autonomous developmental processes not affected by the moment of birth) on macronutrient composition of very preterm milk were studied. Total N, fat, lactose and carbohydrate concentrations, energy density and 24 h volume were determined in 282 24 h milk samples collected at weekly intervals (days 7–55 of lactation) from seventy-nine women delivering their babies between 25 and 29 weeks of gestation. GA related differences were found for carbohydrate concentration only: carbohydrate concentration was lower with increasing GA. PNA was related to a decrease in total N and an increase in lactose concentration. PMA was not related to milk composition. Our data indicate that PNA strongly influences the development of the composition of very preterm human milk, while GA affects carbohydrate content with a negligible effect on the nutritional value of the milk. We conclude that in accordance with current opinion in paediatrics, human milk is the best source of nutrients even for very preterm (< 30 weeks GA) infants.

Human milk: Preterm infant: Development

Human milk has long been recommended as the ideal nutrient source for full-term neonates, but there is still controversy concerning its suitability for the preterm infant (Lawrence, 1994). Most studies comparing the macronutrient composition of preterm human milk with full-term human milk found a difference in composition, with preterm milk having a higher N content and a higher nutritional value than full-term milk (Atkinson *et al.* 1978, 1980; Gross *et al.* 1980, 1981; Schanler & Oh, 1980; Anderson *et al.* 1981; Guerrini *et al.* 1981; Lemons *et al.* 1982, 1983; Butte *et al.* 1984; Lepage *et al.* 1984; Darwish *et al.* 1989; Dawodu *et al.* 1990). These findings gave rise to the common consent that when a mother gives birth prematurely her milk is more suitable for her child than full-term milk. It is also known that the composition of milk shows considerable differences with the stage of lactation (Hyttén, 1954; Atkinson *et al.* 1978, 1980; Gross *et al.* 1980; Anderson *et al.* 1981; Hibberd *et al.* 1982; Anderson *et al.* 1983; Butte *et al.* 1984; Pierse *et al.* 1988; Jain & Bijlani, 1989; Lawrence, 1994).

The underlying factors producing the differences in

composition and the mechanisms leading to the patterns of change in composition are still unclear. In the present study we have examined the influence of gestational age at delivery (GA) and duration of lactation (postnatal age; PNA) and, when indicated, of post-menstrual age (PMA; GA + PNA) on the changing macronutrient composition of preterm milk in a group of mothers giving birth before 30 weeks of gestation. Mathematically, PMA is the simple addition of GA and PNA, but developmentally PMA is an independent time measure, reflecting endogenously-generated maturational processes from conception onwards (Prechtl, 1984). As the moment of conception usually is uncertain, it is common practice to use PMA for documenting developmental age.

Materials and methods

Milk donors

Milk samples (*n* 311) were obtained at weekly intervals from seventy-nine mothers giving birth before 30 weeks of

Abbreviations: GA, gestational age at delivery; PMA, post-menstrual age; PNA, postnatal age.

* **Corresponding author:** Dr Y. G. H. Maas, fax +31 20 6965 099, email y.g.maas@amc.uva.nl

gestation and who had the intention to breast-feed their infants. GA was determined by the first day of the last menstrual period of the mother. This was confirmed either by an ultrasound examination during early pregnancy or a maturational assessment of the preterm infant with the help of the Dubowitz score (Dubowitz *et al.* 1970). Thirteen mothers gave birth in the 25th or 26th week of gestation, twenty mothers in the 27th week, twenty-one mothers in the 28th week and twenty-five mothers in the 29th week.

The study protocol was approved by the Medical Ethical Committee of the Academic Medical Center, Amsterdam.

Collection of milk samples

The collection of the 24 h samples started as soon as there was sufficient milk to both feed the child and take a 25 ml sample for analysis. Samples were taken for as long as the infant stayed in the hospital and milk production was adequate. Mothers pumped their breasts manually or mechanically, collecting the milk in sterile (deionized) bottles. The number and duration of expressions of milk varied per mother, according to their own habit. All expressions of milk were pooled over 24 h, mixed thoroughly and the volume was measured. All samples were stored at -20° until analysed.

Chemical analysis

Total N concentration (mg/kg) was determined using Kjeldhal analysis (Helrich, 1990). Crude protein was calculated by multiplying Kjeldhal N by 6.38. Fat (g/kg) was determined according to the method of Roesse-Gottlieb (Helrich, 1990). Lactose (g/kg) was determined using an enzymic procedure (Boehringer Mannheim GmbH, 1989). Carbohydrate (g/kg) was calculated as:

$$\text{carbohydrate} = \text{DM} - \text{protein} - \text{fat} - \text{ash},$$

DM being determined as the mass left after rotary evaporation at 102° and ash being determined as the mass left after heating at 550° (Helrich, 1990). Gross total energy content (kJ/kg) was calculated as:

$$\text{energy} = (\text{protein} \times 5.65 + \text{fat} \times 9.25 + \text{carbohydrate} \times 3.95) \times 4.18,$$

with protein, fat and carbohydrate expressed in g/kg milk and the constants expressed in kcal/g; 4.18 being the factor used to convert values to kJ (Anderson *et al.* 1981).

Statistical analysis

To evaluate the effects of GA, PNA or PMA on the macronutrient composition of very premature human milk, unbalanced repeated-measurements analysis of covariance with structured covariance matrices was performed (BMDP 5V; Dixon, 1992). This technique allows for missing data which are estimated implicitly from the available data. Only 24 h samples taken before 56 d of lactation (n 282) were used in the statistical analysis, as the older lactational ages lacked sufficient samples for reliable statistics. PMA being the same as GA + PNA makes it impossible to analyse all

three effects (PNA, PMA and GA) in one model. Thus, analysis of the effects of GA and PNA was first performed for each nutrient and energy density using the following model:

$$\begin{aligned} \text{nutrient or energy density} &= \text{GA} + \text{PNA} \\ &+ 24 \text{ h volume effect}, \end{aligned}$$

and for 24 h volume using the model:

$$24 \text{ h volume} = \text{GA} + \text{PNA}.$$

When a statistically significant ($P < 0.05$) effect of GA was found in this model, PNA was replaced by PMA in order to decide whether the combined effect of GA and PNA could be explained by a single effect of PMA. This would be the case if GA lost its significance in the PMA model. If GA remained significant in this model as well, this was seen as an indication of an independent effect of GA. Time scales of PNA and PMA were both divided into eleven time intervals, for PNA between days 7 and 55 (interval 4–5 d) and for PMA between days 183 and 258 (interval 7 d). GA was used as a between-mother grouping variable with four categories: 25–26 weeks, 27 weeks, 28 weeks and 29 weeks. The 24 h milk volume was used as a time-varying covariate in the nutrient and energy density analysis.

To test the assumptions of the model and to check for outliers, analysis of residuals was performed from the unbalanced repeated measurements analysis. When indicated, a Box-Cox (1964) transformation was applied and the effect of outliers was analysed. When a significant overall effect of GA or time was found, a test of which groups or periods differed was performed by calculating contrasts.

Results

24 h milk volume

Based on the residual analysis this variable was log transformed. The amount of milk produced by the mothers during a 24 h period was not found to be related to GA or PNA (Tables 1–3).

Total nitrogen

Changes in total N were related to PNA; an increase in PNA was associated with a significant decrease in total N content (Tables 1–3). Moreover, total N content was related to 24 h milk volume; an increase in 24 h milk volume of 100 ml was associated with a decrease in total N content of 8 (SE 2) mg.

Fat

Fat content was not found to be related to GA or PNA (Tables 1–3), nor was it found to be affected by 24 h milk volume.

Lactose

Based on the residual analysis this variable was transformed using the exponent 2.5. Lactose content was not found to be

Table 1. Milk volume (24 h) and macronutrient composition of preterm human milk for eleven postnatal periods (range 7–55 d)

(Values are means and standard deviations)

Postnatal period	Postnatal age (d)	n	Volume (ml/24 h)		Total N (g/kg)		Fat (g/kg)		Lactose (g/kg)		Carbohydrate (g/kg)		Energy (kJ/kg)	
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	7–10	36	308	192*	3.2	0.8	35.8	9.1	55.0	6.6	71.9	4.8	3054	421
2	11–14	31	261	201	3.0	0.6	34.9	10.5	55.9	6.1†	71.5	6.2	2988	419
3	15–19	32	391	255	2.6	0.6	34.0	5.7	58.9	5.2	74.3	4.2	2930	263
4	20–23	35	329	253	2.5	0.5	33.9	7.5	57.6	4.7	72.5	4.7	2885	361
5	24–28	38	263	193	2.5	0.5	32.8	5.8‡	57.7	4.9	72.6	5.5‡	2841	274‡
6	29–32	24	391	252	2.2	0.4	35.4	5.0	60.2	4.0	75.3	4.0	2946	255
7	33–37	31	334	218	2.1	0.3	35.4	7.4	60.4	4.0	74.5	4.1	2921	306
8	38–41	16	264	243	2.4	0.5	31.7	8.7	57.1	6.6	70.4	5.5	2749	334
9	42–46	18	454	247	2.1	0.3	35.7	4.4	61.8	3.0	75.0	3.5	2933	195
10	47–50	12	366	281	2.0	0.5	33.3	5.4	59.5	5.1	72.7	5.1	2786	210
11	51–55	9	348	204	2.0	0.3	32.7	6.4	61.9	2.1	72.9	4.2	2774	274
all	7–55	282	329	231§	2.5	0.6	34.3	7.3§	58.2	5.4§	73.0	4.9§	2914	329§

* n 35.

† n 30.

‡ n 37.

§ n 281.

Table 2. Milk volume (24 h) and macronutrient composition of preterm human milk for eleven post-menstrual periods (range 183–258 d)

(Values are means and standard deviations)

Post-menstrual period	Post-menstrual age (weeks)	n	Volume (ml/24 h)		Total N (g/kg)		Fat (g/kg)		Lactose (g/kg)		Carbohydrate (g/kg)		Energy (kJ/kg)	
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
1	26–27	1	450		3.8		39.7		57.0		77.1		3384	
2	27–28	9	277	113*	3.2	0.7	34.4	7.5	55.3	3.8	74.7	4.1	3045	318
3	28–29	17	324	213	2.9	0.5	34.8	7.0	57.7	4.6	74.9	3.0	3012	304
4	29–30	36	310	206	2.9	0.7	36.3	9.3	55.9	7.1	72.4	5.4	3035	439
5	30–31	50	300	217	2.8	0.6	34.1	8.7	57.4	5.8†	72.6	5.9	2937	364
6	31–32	54	315	233	2.5	0.6	34.3	6.1‡	58.4	5.0	73.6	4.5‡	2917	274‡
7	32–33	46	366	268	2.3	0.4	33.7	6.7	58.7	5.0	72.4	5.0	2838	284
8	33–34	32	364	269	2.2	0.4	34.0	6.5	60.3	4.4	73.6	4.7	2865	281
9	34–35	22	319	226	2.2	0.5	33.7	7.3	59.2	5.1	72.3	5.1	2822	327
10	35–36	11	390	238	2.0	0.4	34.3	6.0	60.3	3.8	71.8	3.9	2818	252
11	36–37	4	358	175	2.1	0.3	30.3	4.3	63.2	2.1	74.3	2.7	2714	178
all	26–37	282	329	231§	2.5	0.6	34.3	7.3§	58.2	5.4§	73.0	4.9§	2914	329§

* n 8.

† n 49.

‡ n 53.

§ n 281.

related to GA, but lactose content increased highly significantly with increasing PNA (Tables 1–3). Milk volume also affected lactose content; an increase in 24 h milk volume of 100 ml was associated with an increase in lactose content of 0.14 (SE 0.02) g.

Carbohydrate

Based on the residual analysis this variable was transformed using the exponent 4.5. An effect of GA on total carbohydrate was found, mainly due to differences between carbohydrate content at 28 weeks of GA and both the carbohydrate contents at 25–26 and 27 weeks of GA (Table 4). A trend towards an effect of PNA on total carbohydrate content was found ($P=0.057$; Tables 1–3). Analysis of an effect of PMA was therefore indicated, confirming an independent effect of GA on carbohydrate

while no effect of PMA was found. An increase in 24 h milk volume of 100 ml was associated with an increase in total carbohydrate content of 0.10 (SE 0.02) g.

Energy content

Energy content was not found to be related to GA or PNA (Tables 1–3), nor was it found to be affected by 24 h milk volume.

Discussion

The present study, like others, showed that the composition of very preterm milk changes during lactation (Atkinson *et al.* 1980; Gross *et al.* 1980; Schanler & Oh, 1980; Anderson *et al.* 1981; Pamblanco *et al.* 1986; Beijers *et al.* 1992). To explain the observed time effect we evaluated

Table 3. Statistical significance (*P* values) of the effect of gestational age at delivery (GA), postnatal (PNA) or post-menstrual (PMA) age (if analysed) and 24 h milk volume on total nitrogen, fat, lactose, carbohydrate and energy content of 24 h milk (model a) and of the effect of GA and PNA on 24 h milk volume (model b), from mothers delivering preterm

(Statistical method was an unbalanced repeated-measurements analysis of covariance with structured covariance matrices using the models: (a) variable = GA + PNA (or PMA) + 24 h volume effect, (b) 24 h volume effect = GA + PNA)

Variable	Effect of:		
	GA	PNA (or PMA)	24 h volume
Total N	0.85	<0.0001	<0.0001
Fat	0.25	0.69	0.31
Lactose	0.25	<0.0001	<0.0001
Carbohydrate	0.0065 0.013	0.057 0.71*	<0.0001 <0.0001
Energy	0.45	0.011	0.15
24 h volume	0.95	0.89	–

* Effect of PMA.

the effect of three time variables: PNA, GA and, a novelty, PMA. The major findings of our study are that developmental changes in milk composition are largely determined by PNA, minimally by GA and not at all by PMA. This means that the composition of preterm human milk is not determined by autonomous developmental processes related to the moment of conception, but that the maternal body adapts to the moment of precocious delivery. Milk volume (24 h) itself did not show a dependence on GA, PNA or PMA. But using volume as a time-varying covariate in our analysis we found that total N content decreases while lactose and carbohydrate content increase when 24 h milk volume increases, which is in accordance with the literature (Atkinson *et al.* 1978; Gross *et al.* 1981; Lemons *et al.* 1982; Anderson *et al.* 1983). One should be aware of the limitation of our finding, i.e. no effect of PNA on 24 h milk volume. Factors such as stress of delivering extremely preterm infants, 'high-tech' neonatal intensive care units and the absence of breast-feeding all contributed to this finding.

Statistical analysis

The aim of our study was to get more insight into the patterns underlying the changes in nutrient concentration of preterm human milk, whereas other studies focused only on differences between milk obtained from mothers delivering their babies preterm *v.* term. In general, the statistical analysis of previous studies has been done on mean nutrient values of small numbers of mothers at different postnatal days (Schanler & Oh, 1980; Guerrini *et al.* 1981; Anderson *et al.* 1983; Lemons *et al.* 1983; Lepage *et al.* 1984; Darwish *et al.* 1989; Dawodu *et al.* 1990), thereby ignoring the considerable variability in milk volume and nutrient concentration which exists between and within individual mothers, particularly in preterm mothers (Gross *et al.* 1981; Hibberd *et al.* 1982; Anderson, 1984). To allow correction for intra- and inter-individual variations we collected longitudinal data from a total of seventy-nine mothers and used unbalanced repeated measurement

Table 4. Total carbohydrate and lactose content of preterm human milk for four gestational age at delivery (GA) groups

(Values are means and standard deviations)

Gestational age (weeks)	<i>n</i>	Carbohydrate (g/kg)		Lactose (g/kg)	
		Mean	SD	Mean	SD
25–26	57	74.6	3.6	58.5	4.7
27	75	74.3	5.2	59.2	4.2†
28	70	71.4	4.9*	56.9	7.2
29	79	72.2	5.0	58.3	4.9‡
25–29	281	73.0	4.9	58.2	5.4

* Mean value was significantly different from that for GA of 25–26 weeks ($P < 0.001$) and from that for GA of 27 weeks ($P < 0.02$).

† *n* 76.

‡ *n* 78.

analysis. In this way we were able to correct for inter-individual as well as intra-individual variability. Moreover, in our study, possible effects of 24 h volume on the observed differences in macronutrient concentrations between GA, PNA or PMA groups were controlled for by means of the statistical technique of covariance analysis.

Effect of gestational age

We found that total carbohydrate concentration was lower when the GA was higher (Table 4). When differences in composition of very preterm human milk are found to be related to the GA this indicates that the event of birth interrupts the gestational developmental processes occurring in the mammary gland, with a lasting effect on the composition of the milk produced. We have seen this only for the total carbohydrate content. An explanation for this might be that we only studied a small range of GA, a broader range from week 25 to 36 might have led to a different conclusion.

Taking into account that approximately 800 g/kg carbohydrate in human milk is considered to be lactose, we could have expected an effect of GA for lactose as well. However, such an effect was absent (Table 4), possibly due to the relatively large variation in the lactose contribution to the carbohydrate content in our samples (640–930 g/kg).

Effect of postnatal age

Changes in composition related to PNA indicate that there is a relationship with the lactational processes (for example of the mammary gland) initiated at the moment of birth. Like other studies, we found similar postnatal changes in protein (Atkinson *et al.* 1978, 1980; Gross *et al.* 1980, 1981; Schanler & Oh, 1980; Anderson *et al.* 1981; Lemons *et al.* 1982, 1983; Anderson *et al.* 1983; Butte *et al.* 1984; Lepage *et al.* 1984; Darwish *et al.* 1989; Dawodu *et al.* 1990) and lactose content (Gross *et al.* 1980; Anderson *et al.* 1981; Gross *et al.* 1981; Darwish *et al.* 1989; Dawodu *et al.* 1990). Similar postnatal changes are also seen in full-term milk (Atkinson *et al.* 1978, 1980; Gross *et al.* 1980, 1981; Anderson *et al.* 1981; Guerrini *et al.* 1981; Hibberd *et al.* 1982; Lemons *et al.* 1982, 1983; Anderson *et al.* 1983; Butte *et al.* 1984; Jain & Bijlani, 1989; Dawodu *et al.* 1990;

Lawrence, 1994), indicating that the compositions of both preterm and full-term human milk change in a similar way (Anderson, 1984). In our samples lactose content also changes relative to total carbohydrate content from 760 g/kg by days 7–10 to 800 g/kg by days 29–32 and 850 g/kg by days 51–55 (Table 1), which is in accordance with what has been found in full-term human milk (Coppa *et al.* 1991, 1993).

Effect of post-menstrual age

Dependency of milk composition on PMA would imply that changes in milk composition are in accordance with developmental changes in the fetal–maternal unit. This could mean that the changes in milk composition are in accordance with the nutrient requirements of the infant at various developmental stages. We did not find such a ‘teleological’ relationship for any of the studied nutrients.

In conclusion, postnatal changes dominate the development of the composition of very preterm human milk. GA affects only carbohydrate content, with a minor net effect on the nutritional value of the milk.

Acknowledgements

First, we want to thank all mothers for their milk samples and cooperation throughout the study. We also thank J. A. Boerma from the laboratories of Nutricia, The Netherlands, for the chemical analyses of the samples. We are grateful to J. G. Koppe and R. de Leeuw for their support of this project. Also, we are grateful to the referee of the journal to which we submitted the paper previously, for critically reviewing the manuscript. Y. G. H. M. and J. G. were financially supported by Nutricia, The Netherlands. This report is part of a study in fulfilment of the Degree in Philosophy in Science for Y. G. H. M.

References

- Anderson DM, Williams FH, Merkatz RB, Schulman PK, Kerr DS & Pittard WB (1983) Length of gestation and nutritional composition of human milk. *American Journal of Clinical Nutrition* **37**, 810–814.
- Anderson GH (1984) The effect of prematurity on milk composition and its physiological basis. *Federation Proceedings* **43**, 2438–2442.
- Anderson GH, Atkinson SA & Bryan MH (1981) Energy and macronutrient content of human milk during early lactation from mothers giving birth prematurely and at term. *American Journal of Clinical Nutrition* **34**, 258–265.
- Atkinson SA, Anderson GH & Bryan MH (1980) Human milk: comparison of the nitrogen composition in milk from mothers of premature and full-term infants. *American Journal of Clinical Nutrition* **33**, 811–815.
- Atkinson SA, Bryan MH & Anderson GH (1978) Human milk: difference in nitrogen concentration in milk from mothers of term and premature infants. *Journal of Pediatrics* **93**, 67–69.
- Beijers RJW, Graaf FVD, Schaafsma A & Siemensma AD (1992) Composition of premature breast-milk during lactation: constant digestible protein content (as in full term milk). *Early Human Development* **29**, 351–356.
- Boehringer Mannheim GmbH (1989) *Methods of Biochemical Analysis and Food Analysis: Using Test-Combinations*, pp. 80–83. Mannheim, Germany: Boehringer Mannheim GmbH.
- Box GEP & Cox DR (1964) Analysis of transformations. *Journal of the Royal Statistical Society* **26B**, 211–252.
- Butte NF, Garza C, Johnson CA, Smith EO & Nichols BL (1984) Longitudinal changes in milk composition of mothers delivering preterm and term infants. *Early Human Development* **9**, 153–162.
- Coppa GV, Gabrielli O, Pierani P, Catassi C, Carlucci A & Giorgi PL (1993) Changes in carbohydrate composition in human milk over 4 months of lactation. *Pediatrics* **91**, 637–641.
- Coppa GV, Gabrielli O, Pierani P, Zampini L, Rotolli G, Carlucci A & Giorgi PL (1991) Qualitative and quantitative studies of carbohydrates of human colostrum and mature milk. *Rivista Italiana di Pediatria* **17**, 303–307.
- Darwish AEH, Dakroury AM, El-Feel MS & Nour NM (1989) Comparative study on breast milk of mothers delivering preterm and term infants. Protein, fat and lactose. *Nahrung* **33**, 249–251.
- Dawodu AH, Osibanjo O & Damole IO (1990) Nutrient composition of milk produced by mothers of preterm infants in Nigeria. *East African Medical Journal* **67**, 873–877.
- Dixon WJ (editor) (1992) In *BMDP Statistical Software Manual*. Berkeley, Los Angeles and Oxford: University of California Press.
- Dubowitz LMS, Dubowitz V & Goldberg C (1970) Clinical assessment of gestational age in newborn infants. *Journal of Pediatrics* **77**, 1–10.
- Gross SJ, David RJ, Bauman L & Tomarelli RM (1980) Nutritional composition of milk produced by mothers delivering preterm. *Journal of Pediatrics* **96**, 641–644.
- Gross SJ, Geller J & Tomarelli RM (1981) Composition of breast milk from mothers of preterm infants. *Pediatrics* **68**, 490–493.
- Guerrini P, Bosi G, Chierici R & Fabbri A (1981) Human milk: relationship of fat content with gestational age. *Early Human Development* **5**, 187–194.
- Helrich K (editor) (1990) In *Official Methods of Analysis of the AOAC*, 15th ed. Arlington, VA: Association of Official Analytical Chemists Inc.
- Hibberd CM, Brooke OG, Carter ND, Haug M & Harzer G (1982) Variation in the composition of breast milk during the first 5 weeks of lactation: implications for the feeding of preterm infants. *Archives of Disease in Childhood* **57**, 658–662.
- Hyttén FE (1954) Clinical and chemical studies in human lactation. IV. Trends in milk composition during course of lactation. *British Medical Journal* **1**, 249–253.
- Jain S & Bijlani RL (1989) The significance of some significant features of breast milk. *Indian Journal of Physiology and Pharmacology* **33**, 118–128.
- Lawrence PB (1994) Breast milk. Best source of nutrition for term and preterm infants. *Pediatric Clinics of North America* **41**, 925–941.
- Lemons JA, Moye L, Hall D & Simmons M (1982) Differences in the composition of preterm and term human milk during early lactation. *Pediatric Research* **16**, 113–117.
- Lemons JA, Reyman D & Moye L (1983) Amino acid composition of preterm and term breast milk during early lactation. *Early Human Development* **8**, 323–329.
- Lepage G, Collet S, Bouglé D, Kien LC, Lepage D, Dallaire L, Darling P & Roy CC (1984) The composition of preterm milk in relation to the degree of prematurity. *American Journal of Clinical Nutrition* **40**, 1042–1049.
- Pamblanco M, Ten A & Comin J (1986) Proteins in preterm and term milk from mothers delivering appropriate or

- small-for-gestational age infants. *Early Human Development* **14**, 267–272.
- Pierse P, Van Aerde J & Clandinin MT (1988) Nutritional value of human milk. *Progress in Food and Nutrition Science* **12**, 421–447.
- Prechtl HFR (editor) (1984) *Continuity of Neural Functions from Prenatal to Postnatal Life. Clinics in Developmental Medicine*, vol. 94. Oxford: SIMP/Blackwell.
- Schanler RJ & Oh W (1980) Composition of breast milk obtained from mothers of premature infants as compared to breast milk obtained from donors. *Journal of Pediatrics* **96**, 679–681.

© Nutrition Society 1998
